

	Primary vaccination			Booster	Interval Between Doses	How & When
	1st Dose	2nd Dose	3rd Dose		(To Note: there is a minimum gap you have to wait after a Covid infection to have a vaccine, even if you are due another dose)	
Over 18: Except those who are severely immunosuppressed	✓	✓	✗	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose (4 weeks from positive test)	Book by phone or online or attend a drop-in
Over 18: Severely Immunosuppressed	✓	✓	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose (4 weeks from positive test)	Invited by NHS - can attend a drop-in or book if overdue (using your original invite letter)
16/17: no additional risk factors	✓	✓	✗	✓	12 weeks between all doses (12 weeks from positive test)	Book by phone or online or attend drop-in
16/17: At risk for specific medical conditions (not those severely immunosuppressed)	✓	✓	✗	✓	8 weeks between primary doses; 12 weeks from primary course to booster (4 weeks from positive test)	Book by phone or online or attend drop-in
16/17: Severely Immunosuppressed	✓	✓	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster (4 weeks from positive test)	Invited by NHS - can attend a drop-in or book if overdue (using your original invite letter)
12 to 15: No additional risk factors	✓	✓	✗	✗	12 weeks between primary doses (12 weeks from positive test)	Invited by NHS - can attend a drop-in or book if overdue
12 to 15: Specific medical conditions or household contacts of a person with immunosuppression	✓	✓	✗	✓	8 weeks between primary doses; 12 weeks from primary course to booster (4 weeks from positive test)	Invited by NHS - can attend a drop-in or book if overdue
12 to 15: Severely Immunosuppressed	✓	✓	✓	✓	8 weeks between primary doses; 12 weeks from primary course and booster (4 weeks from positive test)	Invited by NHS - can attend a drop-in or book if overdue (using your original invite letter)
5 -11: No additional risk factors	✗	✗	✗	✗	N/A	N/A
5 to 11: Specific medical conditions or household contact of a person with immunosuppression	✓	✓	✗	✗	8 weeks between primary doses (4 weeks from positive infection)	Invited by NHS (From late January 2022)
5-11: Severely Immunosuppressed	✓	✓	✓	✗	8 weeks between primary doses (4 weeks from positive infection)	Invited by NHS (From late January 2022)