

	Primary Course Doses			Booster Doses		Interval Between Doses* (There is a minimum time you need to wait after a Covid Infection to have a vaccine, even if you are due another dose)	How & When
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>		
Over 75	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses; (4 weeks from first symptoms or positive test)	Invited by NHS – Spring (booster 2) invite will come out via post from mid-March onwards
Elderly Care Home Residents	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses with flexibility; (4 weeks from first symptoms or positive test)	Delivered through care home - Spring (booster 2) will be offered from mid-march onwards
Over 18: Except those who are immunosuppressed	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster dose; (4 weeks from first symptoms or positive test)	If not completed the primary course or had booster 1 yet - Book by phone or online or attend a drop in.
Over 12: Severely Immunosuppressed	✓	✓	✓	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose 12 weeks between booster doses (4 weeks from first symptoms or positive test)	Invited by NHS – Spring (booster 2) invite will come out via post from mid-March onwards
Over 12: Immunosuppressed (but not severely immunosuppressed)	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses; (4 weeks from first symptoms or positive test)	Invited by NHS – Spring (booster 2) invite will come out via post from mid-March onwards
16/17: no additional risk factors	✓	✓	✗	✓	✗	12 weeks between all doses ; (12 weeks from first symptoms or positive test)	Book by phone or online or attend drop-in
16/17: At risk for specific medical conditions (not those immunosuppressed)	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster; (4 weeks from first symptoms or positive test)	Book by phone or online or attend drop-in
12 to 15: No additional risk factors	✓	✓	✗	✗	✗	12 weeks between primary doses; (12 weeks from first symptoms or positive test)	Invited by NHS - can attend a drop-in or book if overdue
12 to 15: Specific medical conditions or household contacts of a person with immunosuppression	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster; (4 weeks from first symptoms or positive test)	Invited by NHS - can attend a drop-in or book if overdue
5 to 11: No additional risk factors	✓	✓	✗	✗	✗	12 weeks between primary doses; (12 weeks from first symptoms or positive test)	Invited by NHS and can reschedule using helpline (From mid-March 2022)
5 to 11: Specific medical conditions or household contact of a person with immunosuppression	✓	✓	✗	✗	✗	8 weeks between primary doses; (4 weeks from first symptoms or positive test)	Invited by NHS - can book or reschedule using helpline
5 to 11: Severely Immunosuppressed	✓	✓	✓	✗	✗	8 weeks between primary doses; (4 weeks from first symptoms or positive test)	Invited by NHS - can book or reschedule using helpline

\*To support operational delivery, the [Green Book](#) sets out additional flexibility for this time to be reduced in some instances – e.g. the time between the first and Spring booster can be reduced from 24 weeks, to a minimum of 12 weeks, where people are co-residents in older adult care homes, living together and having a domiciliary visit, or where an Immunosuppressed individual has had an additional primary dose followed by booster 1 at a later date.