

	Primary Course Doses			Booster Doses		Recommended Interval Between Doses* (please note the operational flexibility for spring booster) (There is also a gap required after a covid infection)	How & When
	1 st	2 nd	3 rd	1 st	2 nd		
Over 75 (or turning 75 by 30 June 2022)	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses; (4 weeks from first symptoms or positive test)	Invited by NHS by post – Spring (booster 2) from Mid-March 22 (as they reach 24 wks)
Care Home Residents	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses; (4 weeks from first symptoms or positive test)	Spring (booster 2) offered through care home from mid-march 22
Over 18: Except those who are immunosuppressed	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster dose; (4 weeks from first symptoms or positive test)	If not completed the primary course or had booster 1 yet - Book by phone or online or attend a drop in.
Over 12: Severely Immunosuppressed	✓	✓	✓	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose 24 weeks between booster doses (4 weeks from first symptoms or positive test)	Invited by NHS – Spring (booster 2) invite by post from April 22
Over 12: Immunosuppressed (but not severely immunosuppressed)	✓	✓	✗	✓	✓	8 weeks between primary doses; 12 weeks from primary course to booster dose; 24 weeks between booster doses; (4 weeks from first symptoms or positive test)	Invited by NHS – Spring (booster 2) invite by post from April 22
16/17: no additional risk factors	✓	✓	✗	✓	✗	12 weeks between all doses ; (12 weeks from first symptoms or positive test)	Book by phone or online or attend drop-in
16/17: At risk for specific medical conditions (not those immunosuppressed)	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster; (4 weeks from first symptoms or positive test)	Book by phone or online or attend drop-in
12 to 15: No additional risk factors	✓	✓	✗	✗	✗	12 weeks between primary doses; (12 weeks from first symptoms or positive test)	Invited by NHS - can attend a drop-in or book if overdue
12 to 15: Specific medical conditions or household contacts of a person with immunosuppression	✓	✓	✗	✓	✗	8 weeks between primary doses; 12 weeks from primary course to booster; (4 weeks from first symptoms or positive test)	Invited by NHS - can attend a drop-in or book if overdue
**5 to 11: No additional risk factors	✓	✓	✗	✗	✗	12 weeks between primary doses; (12 weeks from first symptoms or positive test)	Invited by NHS and can reschedule using helpline (From mid-March 2022)
**5 to 11: Specific medical conditions or household contact of a person with immunosuppression	✓	✓	✗	✗	✗	8 weeks between primary doses; (4 weeks from first symptoms or positive test)	Invited by NHS - can book or reschedule using helpline
**5 to 11: Severely Immunosuppressed	✓	✓	✓	✗	✗	8 weeks between primary doses; (4 weeks from first symptoms or positive test)	Invited by NHS - can book or reschedule using helpline

***Operational flexibility for spring booster:** Intervals may be reduced to no less than 12 weeks in these circumstances: 1) care home residents; eligible adults living together and having a home visit, severely immunosuppressed; eligible individuals who turn up at clinics before 24 weeks (and before their scheduled appointment by blue envelope). Eligible individuals who reach their 24 weeks after July 1st will be brought forward for scheduling to June (as long as they are at 12 weeks for June appointments).

****4 year olds turning 5 between 1/4/22 and 31/8/22** may be invited forward. They must have turned 5 in order to receive the vaccine.